

## **Publications:**

Michel, W.C. and Case, J.F. The effects of a water-soluble petroleum fraction on the behavior of the hydroid coelenterate *Tubularia crocea*. *Mar. Environ. Res.* 13: 161-176, 1984.

Zimmer-Faust, R.K., Tyre, J.E., Michel, W.C. and Case, J.F. Chemical mediation of appetitive feeding in a marine decapod crustacean: The importance of suppression and synergism. *Biol. Bull.* 167: 339-353, 1984.

Zimmer-Faust, R.K., Michel, W.C., Tyre, J.E. and Case, J.F. Chemical induction of feeding in California spiny lobster *Panulirus interruptus* (Randall): Responses to molecular weight fractions of abalone. *J. Chem. Ecol.* 10: 957-971, 1984.

Michel, W.C. Mate recognition by an Antarctic isopod crustaceans. In *Olfaction and Taste IX*, edited by S.D. Roper and J. Atema. New York: New York Academy of Sciences, pp. 494-497, 1988.

Ache, B.W., McClintock, T.S., Schmedel-Jakob, I. and Michel, W.C. Evidence for multiple transduction pathways in lobster olfactory receptor neurons. In: *Olfaction and Taste X* (K Döving, ed.), GCS a.s.: Oslo, pp. 101-105, 1990.

Schmedel-Jakob, I., Michel, W.C., Anderson, P.A.V. and Ache, B.W. Whole cell recording from lobster olfactory receptor cells: Multiple ionic bases for the receptor potential. *Chem. Senses* 15: 397-405, 1990.  
Fadool, D.A., Michel, W.C. and Ache, B.W. Odor sensitivity of cultured lobster (*Panulirus argus*) olfactory receptor neurons. *Tissue Cell* 23: 719-732, 1991.

Michel, W.C., McClintock, T.S. and Ache, B.W. Inhibition of lobster olfactory receptor cells by an odor-activated potassium conductance. *J. Neurophysiol.* 65(3): 446-453, 1991.

Michel, W.C. and Ache, B.W. Cyclic nucleotides mediate an odor-evoked potassium conductance in lobster olfactory receptor cells. *J. Neurosci.* 12: 3979-3984, 1992.

Fadool, D.A., Michel, W.C. and Ache, B.W. Odor sensitivity of cultured lobster olfactory receptor neurons is not dependent on process formation. *J. Exp. Biol.* 174: 215-233, 1993.

Michel, W.C. and Ache, B.W. Integration of opposing odor-evoked currents by lobster olfactory receptor neurons. In *Proceedings of the 11th International Symposium on Olfaction and Taste and of the 27th Japanese Symposium on Taste and Smell*, edited by K. Kurihara, N. Suzuki and H. Ogawa. Tokyo: Springer-Verlag, pp. 790-791, 1993.

Michel, W.C., Trapido-Rosenthal, H.G., Chao, E.T. and Wachowiak, M. Stereoselective detection of amino acids by lobster olfactory receptor neurons. *J. Comp. Physiol. [A]* 171: 750-712, 1993.

Michel, W.C., Kohbara, J. and Capril, J. Amino acid receptor sites in the facial taste system of the sea catfish *Arius felis*. *J. Comp. Physiol. [A]* 172: 129-138, 1993.

Boekhoff, I., Michel, W.C., Breer, H. and Ache, B.W. Single odors differentially stimulate dual second messenger pathways in lobster olfactory receptor cells. *J. Neurosci.* 14:3304-3309, 1994.

Corotto, F.S. and Michel, W.C. A hyperpolarization-activated cation conductance in lobster olfactory receptor neurons. *J. Neurophysiol.* 72:360-365, 1994.

Michel, W.C. and Ache, B.W. Odor-evoked inhibition in primary olfactory receptor neurons. *Chem. Sens.* 19: 11-24, 1994.

Michel, W.C. and Lubomudrov, L.M. Specificity and sensitivity of the olfactory organ of the zebrafish *Danio rerio*. *J. Comp. Physiol. [A]* 177: 191-199, 1995.

Corotto, F.S., Piper, D.R., Chen, N.S. and Michel, W.C. Voltage- and Ca<sup>2+</sup>-gated currents in zebrafish olfactory receptor neurons. *J. Exp. Biol.* 119:1115-1126, 1996.

Michel, W.C. and Derbidge, D.S. Evidence of distinct amino acid and bile salt receptors in the olfactory system of the zebrafish *Danio rerio*. *Brain Res.* 764:179-187, 1997.

Corotto, F.S. and Michel, W.C. Mechanisms of afterhyperpolarization in lobster olfactory receptor neurons. *J. Neurophysiol.* 80:1268-1276, 1998.

Ma, L. and Michel, W.C. Drugs affecting phospholipase C-mediated signal transduction block the olfactory cyclic nucleotide-gated current of adult zebrafish. *J. Neurophysiol.* 79: 1183-1192, 1998.

Lipschitz, D.L. and Michel, W.C. Physiological evidence for the discrimination of L-arginine from structural analogs by the zebrafish olfactory system. *J. Neurophysiol.* 82: 3160-3167, 1999.

Michel, W.C., Steullet, P., Cate, H.S., Burns, C.J., Zhainazarov, A.B. and Derby, C.D. High-resolution functional labeling of vertebrate and invertebrate olfactory receptor neurons using agmatine, a channel permeant cation. *J. Neurosci. Meth.* 90: 143-156, 1999.

Michel, W.C. Cyclic nucleotide-gated channel activation is not required for activity dependent labeling of zebrafish olfactory receptor neurons by amino acids. *Biol. Signals Recept.* 8: 338-347, 1999.

Steullet, P., Cate, H.S., Michel, W.C. and Derby, C.D. Functional units of a compound nose: Aesthetasc sensilla house similar populations of olfactory receptor neurons on the crustacean antennule. *J. Comp. Neurol.* 418: 270-280, 2000.

Lipschitz, D.L. and Michel, W.C. Microvillar olfactory sensory neurons detect amino acid odorants. *Chem. Sens.* 27:277-286, 2002.

Edwards, J.G. and Michel, W.C. Odor-stimulated glutamatergic neurotransmission in the zebrafish olfactory bulb. *J. comp. Neurol.* 454:294-309, 2002.

Michel, W.C., Sanderson, M.J., Olson, J.K. and Lipschitz, D.L. Evidence of a novel transduction cascade mediating detection of polyamines by the zebrafish olfactory system. *J. Exp. Biol.* (submitted).

### **Abstracts:**

Corotto, F.S. and Michel, W.C. Mechanisms of afterhyperpolarization in lobster olfactory receptor neurons. *Soc. Neurosci. Abstr.* 58.3, 1995.

Michel, W.C., Burns, C.J. and Derbidge, D.S. A functional activity map of the zebrafish olfactory organ: A new method for activity dependent mapping of odor sensitivity using an agmatine specific antibody. *Chem. Senses* 21: 642, 1996.

Michel, W.C., Derbidge, D.S., Chen, N.S. and Ikeda, D.K. Evidence for distinct receptors and transduction pathways for amino acid and bile acid stimuli in the zebrafish olfactory system. *Chem. Senses* 21: 642-643, 1996.

Ma, L. and Michel, W.C. Ruthenium red and neomycin block the zebrafish olfactory cyclic nucleotide-gated channel. *Chem. Senses* 22: 740, 1997.

Michel, W.C. and Burns, C.J. Spatial organization of odor-stimulated olfactory receptor neurons in the zebrafish olfactory epithelium. *Chem. Senses* 22: 748, 1997.

Michel, W.C., Steullet, P. and Derby, C.D. Activity dependent labeling of spiny lobster olfactory receptor neurons confirms that aesthetasc sensilla are redundant sampling elements. *Chem. Senses* 22: 748, 1997.

- Eram, M., Brown, H.M. and Michel, W.C. Amino acid immunocytochemical characterization of the rat vomeronasal organ. *Chem. Senses* 22: 678, 1997.
- Michel, W.C. Distribution of amino acid immunoreactivity in the developing and adult olfactory organ of zebrafish. *Chem. Senses* 23: 552, 1998.
- Edwards, J.G. and Michel, W.C. Mapping the distribution of ionotropic glutamate receptors in the olfactory bulb of zebrafish using a channel permeant probe, agmatine (AGB). *Chem. Senses* 24: 534-535, 1999.
- Steullet, P., Cate, H.S., Nego, V., Michel, W.C. and Derby, C.D. Functional mapping of olfactory receptor cells within aesthetasc sensilla of the spiny lobster. *Chem. Senses* 24: 634, 1999.
- Lipschitz, D.L. and Michel, W.C. Guanidinium-based arginine analogs are detected by multiple odorant receptors (OR) in the zebrafish (*Danio rerio*) olfactory system. *Chem Senses* 24: 593, 1999.
- Michel, W.C., Danaceau, J.P. and Lucero, M.T. Odor-simulated labeling of individual olfactory sensory receptor neurons in the squid. *Soc. Neurosci. Abstr.* 29: 54.12, 1999.
- Edwards, J.G., Lipschitz, D.L. and Michel, W.C. Transmission of olfactory information from the epithelium to the bulb occurs via glutamate release in zebrafish. *Chem. Senses* 24: (in press).
- Lipschitz, D.L. and Michel, W.C. Do amino acids stimulate ciliated or microvillar olfactory sensory neurons? *Chem. Senses* 25: 669-670, 2000.
- Lucero, M.T. and Michel, W.C. Olfactory neuron subtypes display unique amino acid profiles but overlapping odor sensitivities. *Chem. Senses* 25: 663-664, 2000.
- Edwards, J.G. and Michel, W.C. Evidence of presynaptic acetylcholine receptors on zebrafish olfactory sensory neurons. *Chem. Senses* 26: 544, 2001.
- Eram, M. and Michel, W.C. Comparison of the amino acid composition of vagal and facial nerve innervated taste buds in the channel catfish. *Soc. Neurosci. vol . 27, Program number #287.10*, 2001.
- Eram, M. and Michel, W.C. Amino acid immunoreactivity of channel catfish peripheral taste system. *Chem. Senses* 26: 564, 2001.
- Edwards, J.G. and Michel, W.C. Pharmacological characterization of ionotropic glutamate receptors in the zebrafish olfactory bulb. *Chem. Senses* 27: A109.
- Edwards, J.G. and Michel, W.C. Ionotropic glutamate receptor activation selectively depletes GABA levels in zebrafish olfactory bulb. *Chem. Senses* 27: A109.
- Rolen, S.H., Michel, W.C. and Caprio, J. Update on polyamines as olfactory stimuli in goldfish. *Chem. Senses* 27: A73-A74.
- Sitthichai, A.A., Michel, W.C. and Lucero, M.T. Metabolic profile and odor responsiveness of squid olfactory neuron subtypes. *Chem. Senses* 27: A74.